

Liquid Fertiliser

Seed Dressing

GRAIN FORTE SE

**BROAD SPECTRUM OF TRACES INCLUDING SELENIUM
COMPLEXED WITH FULVIC 2% AND BIOSTIMULANTS**

**Iron 4%, Zinc 4%, Manganese 3%,
Copper 0.6%, Boron 0.3% with Selenium,
Molybdenum and Cobalt.**

Mineral sulphates for rapid plant absorption.

**A microbe base of natural biostimulants,
making every drop count.**

- Fulvic keeps trace element sulphates available for longer.
- Maximise grain / produce quality and weight.
- Address specific trace element deficiencies in crops, pasture, horticulture, vines, treecrops and turf.
- Biostimulants to optimise plant health and growth.
- Natural uptake & wetting agents enhance spray efficiency.
- Fulvic buffers against 'burning' with foliar applications.
- With a CEC of 1500 and tiny molecule size, fulvic provides natural chelation and improved nutrient uptake.



Make Every Drop Count

Nutrient components of liquid fertilisers account for 45-60% of total volume. Normally most of the remaining volume is just water to solubilise the minerals within.

HumiPLEX liquids are different. They are made with a microbial liquid base containing by-products (metabolites) from beneficial soil microbes... also known as biostimulants. The product is filled with natural plant growth hormones, vitamins and immune enhancers. We also include fulvic, a natural chelation and uptake agent... The combination primes plants to accept nutrition for improved uptake and efficiency... *making every drop count!*

TYPICAL ANALYSIS (w/v)

Sulphur	5.7% as sulphate	Cobalt	0.1% as sulphate
Iron	4.0% as sulphate	Selenium	0.0826% as selenate
Zinc	4.0% as sulphate	Molybdenum	0.03% as molybdate
Manganese	3.0% as sulphate	Fulvic acid	2.1%
Copper	0.58% as sulphate	pH	2.1-2.4
Boron	0.32% as boric acid	Specific Gravity	1.28-1.31 g/mL

Available in 20L, 200L & 1000L containers.

Check label for more detailed application and handling information.

RECOMMENDATIONS

	Rate / Ha	Timing
Seed Dressing	5-8 L/tonne	Apply to seed pre-sowing
Foliar	Apply when a deficiency has been identified by tissue, petiole or SAP analysis	
Cereals	2-5 L	From 3-5 leaf stage through to first node (Zadoks GS 13 to 31)
Seeding (In Furrow)	3-5 L	
Canola	2-5 L	4-6 leaf stage (Mid Rosette) ~ Pre-flowering (Late Rosette-Bolt)
Legumes (Beans/Peas/Lupins)	2-5 L	4-6 leaf (2nd-3rd node with sufficient leaf area) ~ Pre-flowering
Pasture	2-5 L	Spray pasture when sufficient leaf area exists
Lucerne	2-5 L	Apply after hay cut when adequate leaf area exists ~ Pre-flowering
Citrus	3-10 L	Spring Flush ~ 2/3 leaf expansion ~ Summer Flush ~ Autumn Flush
Vines	3-5 L	2 weeks post budburst ~ 4 weeks post budburst ~ Pre-flowering ~ Post-Harvest
Vegetables (Potato/Onion/Carrot)	4-8 L	Apply from full leaf emergence ~ Repeat applications as required (10-14 days)
Turf	5-10 L	
Fertigation	5-15 L	

Rates and timings may change depending on crop and season.

Always consult a LawrieCo area manager or distributor for specific recommendations.

Building wealth from soil with
Next Generation Fertiliser