

ANIMAL HUMIC FULVIC

ANIMAL GRADE HUMIC & FULVIC ACIDS

Include with dairy or beef cattle, pig, poultry, sheep, horse & fish feed, supplements or water.


- Improves feed and supplement digestion.
- Faster animal weight gains.
- Increase dairy feed efficiency & reduce mastitis, animal stress & healing time.
- Odour reduction in animal waste.
- Support general health & vitality of animals, birds & fish.



Poultry: Effects of adding Humate over 5 weeks:

Losses decreased by 50%

Active (live) weight increased by 30%



Above results are from wide scale testing in 1998 at the Severny pedigree poultry breeding state farm near Bratsk, Russia. *Teravita, Humates in poultry and stock farming.*

In the diets of beef cattle, "it was concluded that, as a percentage of humate administered was increased the amount of feed intake was decreased, while still increasing weight gain." *Uni Arizona Animal Sciences, "Effects of humic/fulvic acids in beef cattle finishing diets", C.P McMurphy et al*

"The use of Humic Acids and related products in feed improved gut health for better nutrient utilisation as well as improved the health status by working against pathogens by developing immunity." *Uni Leipzig, Germany, "Humic Acid Substances in Animal Ag", KMS Islam et al.*

TYPICAL ANALYSIS (w/w dry basis)

Water Solubility	90% min	pH	10 - 11
Humic Acid (dry basis)	60% min	Size Powder	20 mesh
Fulvic Content	15% min	Colour	Black

RECOMMENDATIONS

ADD to:	Feed rations, drinking water, mineral supplements
Feed rations	1-2kg per tonne of feed
Drinking water	1 gram per litre of water
Mineral supplements	10% by weight of supplementation
Unhealthy Animal, Birds, Fish	Initial support - 20 gram per 100kg of body weight
	Subsequent - 5 gram per 100 kg of body weight

Rates and timings may change depending on crop and season. Always consult a LawrieCo consultant or distributor for specific recommendations.

Building wealth from soil with
Next Generation Fertiliser

