

# BiMAX

## 12% PIRANHA KELP

### 12% AUSTRALIAN BULL KELP AND NORTH ATLANTIC ASCOPHYLLUM NODOSUM

An indispensable biostimulant containing potent chelating agents, natural plant hormones and complex sugars which stimulate and feed plant-growth promoting bacteria, increase resistance to frost and environmental stresses, and improve flowering and fruit set.

20L  
200L  
1000L

Batch

This label complies with the Fertiliser Industry Code of Practice for Fertiliser Description and Labelling

*Wealth from Soil*

LawrieCo www.lawrieco.com.au  
Ph 08 8260 1134 47 Naweena Road, Regency Park SA 5010

#### RECOMMENDATIONS

	Rate / Ha	Timing
<b>Seed Treatment</b>	10L/T seed	Increase root length, mass, shoot growth
Foliar	<b>Additional applications can be made prior to or following stress periods such as frost or drought</b>	
Cereals	3-5 L	Tillering (GS 20-29) ~ stem elongation (GS 30-31) ~ Booting (GS 40-49)
Canola	3-5 L	4-6 leaf stage (Mid Rosette) ~ repeat as required
Legumes (Beans/Peas/Lupins)	3-5 L	4-6 leaf (2nd-3rd node with sufficient leaf area) ~ Pre-flowering
Pasture	3-8 L	Spray pasture when sufficient leaf area exists
Citrus	3-10 L	Early vegetative, 4 week intervals
Vines	3-10 L	4 weeks post budburst ~ pre flowering ~ berries peaseize
Potatoes	6-8 L	Full leaf emergence ~ Tuber initiation ~ 12-14 days later ~ flowering ~ bulking
Onions/Carrots	6-8 L	2-3 weeks after emergence ~ root enlargement ~ 10-14 day intervals until harvest
Turf	10-25 L	3-4 week intervals from initial growth stage ~ after heavy use ~ late season
<b>Fertiligation</b>	10-30 L	

Rates and timings may change depending on crop, season, soil type and fertility.  
Always consult a LawrieCo area manager or distributor for specific recommendations.

#### APPLICATION

##### AGITATE WELL BEFORE USE

**Conditions of use:** Ensure adequate leaf area index for nutrient absorption. Not for use when plant is under stress. e.g. moisture, frost, disease or pests. Spray at moderate ambient temperatures, 11-28 degrees. High humidity will stimulate plant uptake. For best results use early morning or late afternoon. Do not spray in winds > 24 km/h, ideally not over 8 km/h

**Equipment:** Ensure spray equipment is clean, remove undissolved particles that may block nozzles and chemical / disease contaminants prior to use.

Water rate 50-100L/ha • 32-50 mesh filtration • Dilution rate 1:10 • Nozzle size 3-5

**Compatibility:** Physically compatible with a range of nutrients and chemicals. A compatibility jar test is recommended before use, incompatibility is indicated by precipitation and/or settling. Contact LawrieCo / your distributor for further compatibility information

**STORAGE:** Containers must be clearly labelled. Store away from strong oxidisers, strong acids and out of direct sunlight.

**DISPOSAL:** Dispose of container to an approved waste disposal plant. Waste product can be added to any soil or into compost/green waste for disposal.

**GENERAL:** If medical advice is needed, have the product container or label on hand.

**SAFETY DIRECTIONS** Wash hands and exposed skin thoroughly after handling. Wear eye protection.

Dispose of contents/container to an approved waste disposal plant in accordance with local regulations.

##### FIRST AID RESPONSE

**If Inhaled:** Remove victim to fresh air and keep at rest in a position comfortable for breathing. If symptoms persist seek medical advice.

**If Swallowed:** (do NOT induce vomiting) rinse mouth and give plenty of water to drink. Contact a doctor if any symptoms arise.

**If in Eyes:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists, get medical attention

**If skin or hair contact occurs:** Wash with plenty of soap and running water. If skin irritation occurs get medical attention.

If any symptoms persist get medical advice/attention or contact a Poisons Information Centre (phone 13 11 26)

Additional info refer to Safety Data Sheet (SDS Code: 8069)

**Impurities:** Cadmium < 0.05 mg/kg; Lead < 0.03 mg/kg; Mercury < 0.01 mg/kg

**Warning** Do not swallow. The spray from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin.

**Conditions of Sale** The Manufacturer/Distributor warrants that this product contains the ingredients specified, and is fit for the purposes stated on this label, when it is used in accordance with directions, under normal conditions. Only the Manufacturer/Distributor is authorised to make any warranty, guarantee or recommendations concerning the product as the location, rate of application and other conditions are

#### TYPICAL ANALYSIS (w/w)

Nitrogen	0.09% as ammonium	Magnesium	0.095% as organic
Potassium	3.55% total	Manganese	0.015% as organic
	3.05% as carbonate	Iodine	0.0015 as organic
	0.5% as organic	Durvillaea potaorum &	
Sulphur	2.085% total	Ascophyllum nodosum	8.0% total
	2.02% as sulphate	pH	9.0-9.5
	0.065% as organic	Total Solids	12.0%
Calcium	0.125% as organic	Specific Gravity	1.08-1.10